



Case Report

Clinical Study and Analysis of 700 Cases of Pneumonia in Children

Hui We¹, Fang qu¹, Lei sun¹, Yajie lei¹ and Rui wang²

Abstract

Introduction: Pneumonia is a common acute disease of the respiratory tract in childhood. In this paper, 700 cases of pneumonia in Changan Hospital of Xi'an city were summarized. All seasons are easy to occur, especially in winter and spring. It is easy to recurrent attacks affect the growth and development of children, if the treatment is not complete. The clinical data of the children are reported as follows:

Background: From January 2016 to December 2016, 700 cases of children pneumonia in Changan hospital in Xi'an were investigated in 1 pairs of 1. Among the 700 children, 437 were boys and 263 were women; 252 cases aged <1 were took up 36%, 287 cases between the age of 1 and 2 were made up 41%, and 161 cases aged >2 were occupied 23%.

Accessory Examination: In 700 cases of chest radiography, 23 cases of lobar pneumonia accounted for 3.28% of the total. 87 cases of double pulmonary bronchitis were 12.42%, 488 cases of bronchitis were 69.71%, 11 cases of interstitial pneumonia were 15.71%, and 4 cases of lobule in chest. 198 cases of WBC ≥ 10 were 28.28%, 76 cases of neutrophils ≥ 70 were 10.86%, 402 cases of lymphoid ≥ 40 were 57.43%, 267 cases of mononuclear ≥ 10 were 38.14%. 196 cases of syncytial virus positive were took up 28%, 4 cases of adenovirus positive were made up 0.57%, 3 cases of influenza virus positive were occupied 0.43%.

Complication: 2 cases of combined respiratory failure accounted for 0.28%, 13 cases of combined enteritis were 1.85%, 8 cases of combined myocardial injury were 1.14%, 1 case of combined heart failure were 0.14%, 1 case of combined Anemia were 0.14%, 2 cases of combined laryngitis were 0.28%.

Treatment: 368 cases of the second generation cephalosporin's +antivirus were 52.57%, 183 cases of the third generation cephalosporin's + antivirus were 26.14%, 129 cases of Cephalosporin +Archie were 18.43%, 20 cases of Penicillin + antiviral were 2.85%, 23 cases of >14 days of treatment were 3.28%, 72 cases of 7-10 days of treatment were 10.28%, 545 cases of 5-7 days of treatment were 77.86%, 60 cases of <5 days of treatment were 8.57%.

Result: Treatment, Infantile pneumonia is the main anti infection and treat the complications. At the time of anti-infection, a wide range of antimicrobial spectrum, less drug resistant bacteria, and low toxicity of liver and kidney are used, combined use of two kinds of broad-spectrum antibiotics, Such as cephalosporin's, macrolides,

and antiviral drugs, and give oxygen therapy, nutritional support, Correcting the acid-base balance and electrolyte disturbance in some children, actively dealing with the complications and avoiding heart failure.

Keywords

Treatment; Injury; Children; Pneumonia; Complications

Case Report

According to the latest research reports, Pneumonia is still the first cause of the death of the in our country, Serious threat to children's health, Because of its serious clinical manifestations, rapid progress, and a dangerous condition, the mortality rate has remained high. There is no definite diagnostic standard, and the diagnosis of etiology is very difficult. Through the analysis of the results of 700 cases of pneumonia, children's pneumonia is not only caused by some bacteria or viruses, but also often caused by the mixed infection of both. Some children have mycoplasma infection. Through the above analysis, we can draw a conclusion (Table 1). The cure rate of Cephalosporin antiviral drugs is 90%, the cure rate of cephalosporin and Archie is over 90%, it is shows that Children pneumonia is an infectious lung disease caused by a variety of pathogenic microbes and protozoa [1-8]. Because of low immune function in children and imperfect defense function of respiratory system, pediatric pneumonia has become one of the most common diseases in children. Morbidity and mortality are far higher than those in young adults. The incidence of pneumonia in children is rapid, respiratory symptoms are obvious, and the condition of the disease is progressed rapidly, and other complications can occur. If the treatment is unseasonal, it can lead to serious consequences. The clinical manifestations of partial pneumonia in children are not typical, which results in the early diagnosis of pneumonia in children, and it is easy to develop into severe pneumonia. At present, children with pneumonia disease are more serious, there will be more factors to induce the emergence of pneumonia, seriously infringing the health of the children's lungs. This is some of the knowledge that we have to know. The clinical manifestations of respiratory symptoms is shortness of breath, cough, expectoration, mostly white phlegm, Other manifestations were fever, drooping spirit loss of appetite, abdominal pain, diarrhea, vomiting. The lungs can be heard dry and wet rales, individual can hear wheeze sounds. The necessary examination he showed that the total number of white blood cells increased and the classification of neutrophils increased. The chest X-ray, or CT, showed that the double lung texture was thickened, and the increase was blurred, visible scattered in between, high density shadow, edge blurred. Part of the children was large, reticulate. The common pathogenic bacteria of bacterial pneumonia are Streptococcus pneumonia, Staphylococcus aureus, Streptococcus, gram-negative tumors and so on. Bronchopneumonia caused by Streptococcus pneumonia is the most common in infancy. The common virus of viral pneumonia is respiratory syncytial virus, adenovirus, influenza virus and Para-influenza virus; In addition, there are mycoplasma, chlamydia, fungi, protozoa, etc. Pathogens are mostly invaded by respiratory tract, causing bronchioles, alveoli and interstitial lung inflammation, causing small airway stenosis or even obstruction, resulting in ventilation disorders. Inflammation causes

*Corresponding author: Rui wang, Xi'an Medical University, Department of Clinical Pediatrics, China, E-mail: jiaofy@yeah.net

Received: March 26, 2018 Accepted: April 11, 2018 Published: April 25, 2018

Table 1: Healed children are automatically discharged from the hospital.

Curative effect	Cephalosporin + Antiviral	Cefotaxime + Archie	Penicillin + Antivirus
Cure	503 people 71.85%	102 people 14.57%	15 people 2.14%
Improve	35 people 5%	22 people 3.14%	4 people 0.57%
Healed	13 people 1.86%	5 people 0.71%	1 people 0.14%

thickening of respiratory membranes and inflammatory exudates in alveolar cavity, resulting in ventilation disorder. Hypoxic and carbon dioxide retention, even respiratory failure caused by ventilation and ventilation disorder, the reflex contraction of the pulmonary artery causes increased pulmonary arterial pressure, and the effect of pathogens and toxins can cause toxic myocarditis. The results of two factors together lead to psychological exhaustion, hypoxia and the role of pathogen toxin. It can also cause toxic encephalopathy, hemorrhage of digestive tract and toxic intestinal paralysis, as well as metabolic and respiratory acidosis, electrolyte disorders encephalopathy, disseminated intravascular coagulation may occur in severe cases.

Conclusion

In the developing countries, Lower respiratory tract infection is usually the main cause of death or next to diarrhea, The most common cause of the disease is bacteria, of which Streptococcus pneumonia is the most common, other pathogens including anaerobic bacteria, Staphylococcus aureus, Homophiles influenza, parrot fever, Chlamydia trachomatis, M. Mora and Legionella pneumophila, Klebsiella pneumonia and other gram negative bacillus coli. Mycoplasma pneumonia (MP) is a particularly common cause of older children and young people, common in spring. The main pathogens of pneumonia in infants and children are viruses: respiratory syncytial virus, adenovirus, Para-influenza virus, type A and B influenza virus [9]. With the widespread use of antibiotics and the changes in the body's responsiveness, low immune function in children and imperfect defense function of respiratory system, pediatric pneumonia has become one of the most common diseases in children. Morbidity and mortality are far higher than those in young adults. The incidence of pneumonia in children is rapid, respiratory symptoms are obvious, and the condition of the disease is progressed rapidly, and other complications can occur. Infantile pneumonia is the main anti infection and treat the complications. At the time of anti-infection, a wide range of antimicrobial spectrum, less drug resistant bacteria, and low toxicity of liver and kidney are used, combined use of two kinds of broad-spectrum antibiotics, Such as cephalosporin's, macrolides, and antiviral drugs, and give oxygen therapy, nutritional support, Correcting the acid-base balance and electrolyte disturbance in some children, actively dealing with the complications.

Reference

- Nir A, Nasser N (2005) Clinical value of NT-ProBNP and BNP in pediatric cardiology. J Card Fail 11: S76-S80.
- Koulouri S, Acherman RJ, Wong PC (2004) Utility of β type natriuretic peptide in differentiating congestive heart failure from lung disease in pediatric patients with respiratory distress. Pediatr Cardiol 25: 341-346.
- Marcelo AP (2000) Pathophysiology of pediatric heart failuer. Pediatric Cardiol 11: 175-184.
- Abdula R (2004) Drug therapy in pediatric cardiology and the role of digoxin in pediatric congeative heart failure. Pediatr Cardiol 25: 621.
- Barnes S, Shiel.Ds B, Bonney W (2004) The pediatric cardiology phartnacopoeia; 2004 update. Pediatr Cardiol 25: 623-646.

- Enwere G, Cheung YB, Zaman SMA (2007) Epidemiology and clinical features of pneunonia according to radiographic findings in Gambian children. Tropical Med Int Health 12: 1377-1385.
- WalRer CF, Black RE (2004) Zinc and the risk for infectiones disease. Annu Rev Nutr 24: 255-275.
- Bhatnagar S, Natchu UC (2004) Zinc in child health and disease. Indian J Pediatr 71: 991-995.
- Booker PD (2002) Pharmacological support for children with myocardial dysfunction. Pediatr Anaesth 12: 5-25.

Author Affiliations

[Top](#)

¹Dept of Pediatrics, Changan Hospital, Xi'an, China

²Xi'an Medical University, Department of Clinical Pediatrics, China