

# Pregnancy Outcomes in Women with Extremely High sFlt-1/PlGF Ratio: Case Series

Oshovskyy V<sup>1</sup>, Arbuzova S<sup>2</sup>, Nikolenko M<sup>3</sup> and Poliakova Y<sup>4\*</sup>

<sup>1</sup>Uniklinika Medical Centre, Kyiv, Ukraine

<sup>2</sup>Professor, Kyiv, Ukraine

<sup>3</sup>Medico-Biological Center "GENOM", Kyiv, Ukraine

<sup>4</sup>Multidisciplinary Clinic of the St. Nicholas, Zaporizhzhia, Ukraine

## Abstract

**Background:** Preeclampsia (PE) is a multisystem disorder, usually defined as the development of hypertension and proteinuria after 20 weeks of pregnancy. The sFlt-1/PlGF ratio has been widely studied as a diagnostic and prognostic marker of preeclampsia and other manifestations of placental dysfunction. A sFlt-1/PlGF ratio greater than 85 for early PE, <34 weeks of gestation suggests a high risk of PE requiring close clinical monitoring.

**Objective:** To evaluate the maternal and perinatal outcomes of pregnancies with an extremely high sFlt-1/PlGF ratio.

**Case Presentation:** The analysis included data on placental growth factor and soluble fms-like tyrosine kinase serum levels, measured during 2017-2020 in 128 pregnant women. Here we present 8 cases of women with a numerical ratio greater than 850.

In all 100% of cases, the signs of obstetric angiogenic catastrophe requiring imminent delivery developed soon.

**Conclusion:** We observed a trend for worsening perinatal outcomes in women with an extremely high sFlt-1/PlGF of  $\geq 850$ .

**Keywords:** Placental Dysfunction; Preeclampsia; sFlt-1/PlGF Ratio

\***Correspondence to:** Poliakova Yevheniia, Multidisciplinary Clinic of the St. Nicholas, Zaporizhzhia, Ukraine; E-mail: [jenyapolyakova@gmail.com](mailto:jenyapolyakova@gmail.com)

**Citation:** Oshovskyy V, Arbuzova S, Nikolenko M, Poliakova Y (2021) Pregnancy Outcomes in Women with Extremely High sFlt-1/PlGF Ratio: Case Series. *J Womens Health Care Manage*, Volume 2:2. 117. DOI: <https://doi.org/10.47275/2692-0948-117>

**Received:** December 28, 2020; **Accepted:** January 20, 2021; **Published:** January 25, 2021

## Introduction

Pregnancy becomes a life-threatening condition for 10 million women annually. The most common complication is preeclampsia, which complicates 5 to 10% of pregnancies in the United States and up to 18% in some regions of Africa [1].

Preeclampsia is the leading cause of death for young women, particularly those living in resource-limited settings, and it leads to approx. 76,000 maternal deaths and 500,000 infant deaths worldwide each year [1].

Preeclampsia (PE) is a multisystem disorder, usually defined as the development of hypertension and proteinuria after 20 weeks of pregnancy [2]. Recent PE pathophysiology studies open up interesting prospects for screening for this disease.

As a sign of placental dysfunction, preeclampsia is associated with dysregulation of pro- and antiangiogenic factors [3]. Although the etiology of PE has not been fully understood, it is believed to occur due to impaired placentation in early pregnancy. Defective trophoblast invasion and abnormal remodeling of spiral arteries result in the

placental hypoperfusion and, as a consequence, leads to oxidative stress [4]. Therefore, placenta releases a large number of antiangiogenic factors into the maternal circulation [5-7]. During normal pregnancy, concentrations of antiangiogenic factor sFlt-1 (soluble fms-like tyrosine kinase-1) remains low, allowing accurate transmission of signals induced by proangiogenic factors such as VEGF (vascular endothelial growth factor) and PlGF (placental growth factor). This balance is crucial to maintain physiological vasodilation [8]. Secondary to hypoperfusion, placenta increases the synthesis of sFlt-1, trying to raise maternal blood pressure and increase placental perfusion. Consequently, levels of circulating proangiogenic factors are reduced and vascular homeostasis is altered, causing endothelial dysfunction, which leads to hypertension, proteinuria and other multiorgan manifestations of PE in mother [9].

Measurements of blood pressure and proteinuria are considered the gold standard for the diagnosis of preeclampsia, although they do not predict the adverse perinatal outcomes with adequate accuracy [10]. Clinically, hypertension and proteinuria do not always reflect the severity of the disease.



The sFlt-1/PlGF ratio has been widely studied since 2004 as a diagnostic and prognostic marker of preeclampsia and other manifestations of placental dysfunction [11]. To date, a large prospective clinical trial PROGNOSIS has clearly demonstrated that the sFlt-1/PlGF ratio  $\leq 38$  may be used to rule out PE within one week, independently of gestational age [12]. A sFlt-1/PlGF ratio greater than 85 (for early PE,  $<34$  weeks of gestation) or 110 (for late PE,  $\geq 34$  weeks) suggests a high risk of PE requiring close clinical monitoring [13].

Thus, estimated sFlt-1/PlGF ratio can be of great added value when used as an add-on to routine assessment methods. Accurate prediction of PE is important, since severe symptoms may worsen both maternal and fetal status.

The main objective of the expectant management of PE is to improve perinatal outcomes owing to delayed delivery, thereby reducing neonatal morbidity and mortality. In 2012, Verlohren S, et al. (2014) [14], reported that a sFlt-1/PlGF ratio above 655 identified women at high risk of imminent preterm delivery before 34 weeks of gestation, who therefore required close monitoring and medical care.

However, in a study by Stolz M, et al. (2018) [2], a sFlt-1/PlGF ratio above 655 did not predict worsening of perinatal outcomes and was not reliable enough to predict results in cases with clinical signs of PE. Nevertheless, the data obtained suggest that an extremely high sFlt-1/PlGF ratio greater than 850 may be more helpful.

## Objective

Our main aim was to evaluate the maternal and perinatal outcomes of pregnancies with an extremely high sFlt-1/PlGF ratio.

## Materials and Methods

The analysis included data on placental growth factor (PlGF) and soluble fms-like tyrosine kinase (sFlt-1) serum levels, measured during 2017-2020 in 128 pregnant women at 18-39 weeks of gestation at the medical centers Genome and Unclinic, Kyiv, Ukraine. The analysis was performed using the analytical method TRACE (Time Resolved Amplified Cryptate Emission) and test systems Thermo Fisher Scientific by BRAHMS Kryptor (Germany). The study was performed on an automatic BRAHMS Kryptor Compact Plus analyzer in full accordance with the manuals for the test kits. The variations in the concentrations of the studied analytes in the examined sample were 8 to 7,100 pg/ml (PlGF), 70 to 68,000 pg/ml (sFlt-1). These results were used to estimate the sFlt-1/PlGF ratio. Patients with a numerical ratio greater than 850 (exposure factor selected based on the literature) were

included in the group for further research. Here we present 8 cases with an extremely high sFlt-1/PlGF ratio. All patients gave consent for the use of their data in a blinded manner.

## Results

Detailed clinical profiles of patients are presented in Table 1. The mean age of pregnant women was  $28.4 \pm 4$  years. The mean gestational age was  $26 \pm 5$  weeks throughout the study. The majority of pregnant women had signs of intrauterine growth restriction (IUGR) (62.5%) and/or preeclampsia (50%) throughout the study.

The mean sFlt-1/PlGF ratio was 1,747 pg/ml.

In all 100% of cases, the signs of obstetric angiogenic catastrophe requiring imminent delivery developed soon. The mean time to delivery was 10 days. In 50% of pregnant women, urgent delivery was required within 48 hours after analysis. Favorable neonatal outcomes were reported in 37.5% of women.

## Discussion

Biochemical and biophysical parameters, routinely used to diagnose PE and determine its severity, such as proteinuria, blood pressure, transaminases, platelets, and serum creatinine, are of limited value in predicting adverse maternal and perinatal outcomes [9]. Finding parameters to stratify patients based on the severity of preeclampsia is a global problem. Except for those patients who are admitted to hospital with clinical signs of severe preeclampsia and/or imminent eclampsia (headache, abdominal pain, uncontrolled blood pressure), it is very difficult to assess the severity of the disease at the first examination [4].

The sFlt-1/PlGF ratio allows clinicians to predict the onset of PE not only in high risk but also in moderate risk patients and to stratify its level, in particular, by confirming or refuting threatening angiogenic profile [8].

In their study, Verlohren S, et al. (2014) [10], reported that the sFlt-1/PlGF ratio was a reliable tool for identifying women at high risk of imminent iatrogenic delivery, demonstrating that the time to delivery was significantly shorter in women with a ratio above 655 [10].

However, recently Stolz M, et al. (2018) [2], compared perinatal outcomes in a group of 30 patients with preeclampsia and sFlt-1/PlGF ratio  $>655$  and 30 patients with preeclampsia and sFlt-1/PlGF ratio  $<655$  [2]. No significant differences in perinatal outcomes have been observed between the groups in this small study.

**Table 1:** Baseline clinical characteristics of patients.

	Maternal age	Gestational age at enrolment	Preeclampsia	IUGR	sFlt-1/PlGF ratio	Time to delivery (days)	Indication for delivery	Adverse perinatal outcomes
1	30	24	Yes BP 160/100	Yes	1631	21	Placenta abruption	Early neonatal mortality
2	34	21	No	Yes	885	2	Severe PE	Intrauterine demise
3	24	29	Yes BP 170/110	No	2048	1	Eclampsia	No
4	29	34	Yes BP 140/100	No	1152	14	Eclampsia	No
5	34	32	Yes BP 170/100	No	2312	1	Placenta abruption	No
6	26	21	No	Yes	2888	2	Severe PE	Early neonatal mortality
7	23	23	No	Yes	1992	7	Severe PE	Early neonatal mortality
8	27	26	No	Yes	1069	34	Severe PE	Intrauterine demise



Currently, very little is known about the significance of exceeding the cut-off of 655 during pregnancy complicated by PE. Even less is known about pregnancies without preeclampsia, but with an extremely high sFlt-1/PIGF ratio. When placental dysfunction occurs in early pregnancy, expectative strategy or preventive intake of acetylsalicylic acid are routinely used, but the course of the disease is often difficult to predict.

In our case series, perinatal outcomes in women with signs of placental dysfunction and an extremely high sFlt-1/PIGF ratio have been analyzed. We sought to further demonstrate that the sFlt-1/PIGF ratio may be used to predict perinatal outcomes. This would determine the cut-off for iatrogenically induced delivery or justify expectative clinical strategy in relevant cases.

In our case series, 100% of women with an extremely high sFlt-1/PIGF ratio developed signs of obstetric catastrophe and required imminent delivery. We suggest that the extremely high sFlt-1/PIGF ratio (over 850 in our study) should be considered as a criterion for inpatient monitoring of mother and fetus to prevent obstetric catastrophes.

## Conclusions

The sFlt-1/PIGF ratio  $\geq 655$  does not appear to be a reliable cut-off for predicting adverse perinatal outcomes in women with clinical signs of preeclampsia and therefore has limited value for stratification of risk groups. However, we observed a trend for worsening perinatal outcomes in women with an extremely high sFlt-1/PIGF  $\geq 850$ .

Our results also confirm that when an extremely high sFlt-1/PIGF ratio is detected, the pregnant woman should be closely monitored and the need for corticosteroids to accelerate fetal lung maturation should be considered.

## Prospects for Future Studies

Further studies are needed to confirm the prognostic role of sFlt-1/PIGF ratio, to determine the cut-off value, and to improve the clinical evaluation of patients with extremely high sFlt-1/PIGF ratios.

## Declaration of Interests

### Ethical Approval

Not applicable.

### Conflicting Interests

The Authors declare that there is no conflict of interest.

### Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

### Informed consent

Written informed consent was obtained from the patients for anonymized information to be published in this article.

## Author Contributions

Material preparation, data collection and analysis were performed by Oshovskyy Viktor and Poliakova Yevheniia.

The first draft of the manuscript was written by Oshovskyy Viktor and all authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

## References

- Robertson SA (2019) Preventing preeclampsia by silencing soluble Flt-1?. *N Engl J Med* 380: 1080-1082. <https://doi.org/10.1056/NEJMcibr1817501>
- Stolz M, Zeisler H, Heinzl F, Binder J, Farr A (2018) An sFlt-1:PIGF ratio of 655 is not a reliable cut-off value for predicting perinatal outcomes in women with preeclampsia. *Pregnancy Hypertens* 11: 54-60. <https://doi.org/10.1016/j.preghy.2018.01.001>
- Herraiz I, Simón E, Gómez-Arriaga PI, Martínez-Moratalla JM, García-Burguillo A, et al. (2015) Angiogenesis-related biomarkers (sFlt-1/PLGF) in the prediction and diagnosis of placental dysfunction: an approach for clinical integration. *Int J Mol Sci* 16: 19009-19026. <https://doi.org/10.3390/ijms160819009>
- De Oliveira L, Peraçoli JC, Peraçoli MT, Korkeas H, Zampieri G, et al. (2013) sFlt-1/PIGF ratio as a prognostic marker of adverse outcomes in women with early-onset preeclampsia. *Pregnancy Hypertens* 3: 191-195. <https://doi.org/10.1016/j.preghy.2013.02.003>
- Tsatsaris V, Fournier T, Winer N (2010) Physiopathologie de la prééclampsie. *Ann Fr Anesth Reanim* 29: e13-e18. <https://doi.org/10.1016/J.ANNFAR.2010.02.011>
- Chaiworapongsa T, Chaemsaihong P, Yeo L, Romero R (2014) Pre-eclampsia part 1: current understanding of its pathophysiology. *Nat Rev Nephrol* 10: 466-480. <https://doi.org/10.1038/nrneph.2014.102>
- Wang A, Rana S, Karumanchi SA (2009) Preeclampsia: the role of angiogenic factors in its pathogenesis. *Physiology* 24: 147-158. <https://doi.org/10.1152/physiol.00043.2008>
- Tardif C, Dumontet E, Caillon H, Misbert E, Dochez V, et al. (2018) Angiogenic factors sFlt-1 and PlGF in preeclampsia: Prediction of risk and prognosis in a high-risk obstetric population. *J Gynecol Obstet Hum Reprod* 47: 17-21. <https://doi.org/10.1016/j.jogoh.2017.10.007>
- Simón E, Permy C, Sacristán L, Zamoro-Lorenci MJ, Villalain C, et al. (2020) sFlt-1/PIGF ratio for the prediction of delivery within 48 hours and adverse outcomes in expectantly managed early-onset preeclampsia. *Pregnancy Hypertens* 22: 17-23. <https://doi.org/10.1016/j.preghy.2020.07.007>
- Verlohren S, Herraiz I, Lapaire O, Schlemmbach D, Zeisler H, et al. (2014) New gestational phase-specific cutoff values for the use of the soluble fms-like tyrosine kinase-1/placental growth factor ratio as a diagnostic test for preeclampsia. *Hypertension* 63: 346-352. <https://doi.org/10.1161/HYPERTENSIONAHA.113.01787>
- Chang YS, Chen CN, Jeng SF, Su YN, Chen CY, et al. (2017) The sFlt-1/PIGF ratio as a predictor for poor pregnancy and neonatal outcomes. *Pediatr Neonatol* 58: 529-533. <https://doi.org/10.1016/j.pedneo.2016.10.005>
- Zeisler H, Llorba E, Chantraine F, Vatish M, Staff AC, et al. (2016) Predictive value of the sFlt-1:PIGF ratio in women with suspected preeclampsia. *N Engl J Med* 374: 13-22. <https://doi.org/10.1056/NEJMoa1414838>
- Caillon H, Tardif C, Dumontet E, Winer N, Masson D (2018) Evaluation of sFlt-1/PIGF ratio for predicting and improving clinical management of pre-eclampsia: Experience in a specialized perinatal care center. *Ann Lab Med* 38: 95-101. <https://doi.org/10.3343/alm.2018.38.2.95>
- Verlohren S, Herraiz I, Lapaire O, Schlemmbach D, Moertl M, et al. (2012) The sFlt-1/PIGF ratio in different types of hypertensive pregnancy disorders and its prognostic potential in preeclamptic patients. *Am J Obstet Gynecol* 206: 58.e1-58.e8. <https://doi.org/10.1016/j.ajog.2011.07.037>